

Claims

1. A method for at least partially separating nucleic acid molecules in a sample into populations wherein a population is tagged or capable of being tagged with a moiety capable of being immobilised on a matrix, said method comprising contacting the nucleic acid containing sample with a matrix whereby the tagged molecules are captured by the matrix and thereby separated from untagged molecules.
2. A method according to claim 1 wherein a population of molecules is tagged or capable of being tagged with a moiety capable of being immobilised on a matrix either directly or indirectly via a binding partner for the tag, said method comprising contacting the nucleic acid containing sample with a matrix, or, where the tag interacts indirectly with the matrix by means of a binding partner, with a binding partner for the tag and with a matrix, whereby tagged molecules are captured by the matrix and thereby separated from untagged molecules.
3. A method as claimed in any one of the preceding claims wherein the tag is a moiety which can be incorporated into a nucleic acid molecule or a moiety which has an affinity for a nucleic acid molecule.
4. A method as claimed in any one of the preceding claims wherein the tag interacts directly with the matrix.
5. A method as claimed in any one of claims 1 to 3 wherein the tag interacts indirectly with the matrix by means of a binding partner for the tag.
6. A method as claimed in any one of the preceding

09870128 "053004
T0059" 827059

claims wherein the tag is a ligand.

7. A method as claimed in claim 6 wherein the ligand is selected from biotin or fluorescein.

5

8. A method as claimed in any one of claims 1 to 5 wherein the tag is a steroid or a steroid like molecule.

9. A method as claimed in claim 8 wherein the steroid is digoxigenin.

10

10. A method as claimed in any one of claims 1 to 5 wherein the tag is an antigen.

11. A method as claimed in any one of claims 1 to 5 wherein the tag is a protein.

15

12. A method as claimed in claim 11 wherein the protein is a nucleic acid binding protein.

20

13. A method as claimed in any one of claims 1 to 5 wherein the tag is a nucleic acid molecule.

14. A method as claimed in any one of claims 1 to 7 wherein the tag is biotin, and the tag is captured by means of streptavidin or avidin.

25

15. A method as claimed in any one of the preceding claims wherein the matrix is in the form of sheets, gels, filters, membranes, fibres, tubes, microtitre plates, columns, particles.

30

16. A method as claimed in claim 15 wherein the matrix is a porous material.

35

17. A method as claimed in claim 16 wherein the matrix is incorporated into a separation device selected from

09070128 053001

centrifuge vials, microtiter plates, cartridges and syringes.

18. A method as claimed in any one of claims 15 to 17
5 wherein an absorbent pad is located on said porous material, a liquid impermeable sheet is located on the face of said absorbent pad remote from said porous material, and a liquid impermeable sheet having one or more holes therein is located on the face of said porous
10 material remote from said absorbent pad, whereby the test sample is applied to one of said holes and is caused to diffuse transversely through said porous material by absorption into said absorbent pad.
19. A method as claimed in any one of the preceding
15 claims for the at least partial separation of a product of restriction enzyme digestion, or for purifying the products of the PCR reaction.
20. A method as claimed in any one of the preceding
20 claims for at least partially separating a mixture of restriction enzyme digested fragments of DNA wherein the starting material is a linear DNA molecule which is tagged or capable of being tagged at or near one or both
25 ends with a moiety capable of being immobilised on a matrix, said method comprising subjecting the DNA molecule to restriction enzyme digestion followed by contacting the sample with a matrix whereby the tagged molecules which originate from an end of the starting
30 material are captured by the matrix and are thereby separated from untagged molecules.
21. A method as claimed in any one of the preceding
35 claims for at least partially separating the correct and desired products of PCR amplification from PCR products which result from incorrect annealing of a PCR primer to template, wherein the template nucleic acid molecule

09870128 "053004

comprises a unique restriction enzyme recognition site at or towards an end of the template, and a PCR primer which is tagged or capable of being tagged is complementary to a sequence on the template which
5 extends partially into the unique restriction site, the method comprising amplifying the template by means of PCR, digesting the PCR products with the restriction enzyme specific for the said unique restriction enzyme recognition site, and contacting the resulting product
10 with a matrix capable of sequestering the tag whereby tagged nucleic acid molecules are captured by the matrix and thereby separated from untagged molecules.

22. A method as claimed in claim 21 wherein the tag is
15 attached at the 5' end of the primer.

23. A method as claimed in any one of claims 1 to 18 for diagnostic PCR.

24. A method as claimed in claim 23 of diagnostic PCR
20 wherein a test sample is separately subjected to PCR reactions, in which the mutation, if present, is in the sequence to which the 3' primer is complementary, a first PCR reaction using a 3' primer complementary to
25 the normal target nucleic acid and a second PCR reaction using a 3' primer complementary to the mutant target, wherein the 3' nucleotide of the mutant primer corresponds to one of the nucleotides which is mutated in the mutant nucleic acid, each of said 3' primers
30 bearing a tag or being capable of being tagged on the 3' nucleotide of the primer, wherein the presence or absence of the tag in the PCR reaction products is detected.

25. A method as claimed in claim 23 of diagnostic PCR
35 of mutations in a nucleic acid molecule wherein the presence or absence of a mutation in a nucleic acid

09070428 053004
F0050 03202850

sample is detected, wherein a 3' primer specific for either the normal or mutant nucleic acid is used, wherein said primer is complementary to a region of the nucleic acid where there is a base difference between the normal and mutated DNA with the 3' terminus of the primer corresponding to the position in the sample where there is a difference between the normal and mutant, the primer being tagged or capable of being tagged, whereby the presence or absence of the tag in the PCR product is detected.

26. A method as claimed in claim 23 of diagnostic PCR in which a test sample is subjected to PCR with primers complementary to the normal target DNA wherein the primer for extending at the 3' end of the target anneals to a sequence the 3' nucleotide of which is mutated in the mutant, the 3' primer bearing a tag or being capable of being tagged at or on the 3' nucleotide, wherein the presence of the tag in the PCR reaction product as detected.

27. A method as claimed in any one of claims 1 to 18 for separating linear from circular DNA molecules.

28. A method as claimed in claim 27 of separating linear from circular nucleic acid molecules in a sample said method comprising introducing a tag to an end of a linear nucleic acid molecule, wherein said tag is capable of being immobilised on a matrix, by direct interaction with the matrix or by indirect interaction by means of a binding partner to the tag, and contacting the sample with a matrix or, where the tag interacts indirectly with the matrix, with the binding partner to the tag and with a matrix, whereby said tagged linear nucleic acid molecules are immobilised on the matrix.

09070128 053001

29. A method as claimed in any one of claims 1 to 18 for *in vitro* packaging of bacteriophage particles.

5 30. A method as claimed in claim 29 of *in vitro* phage packaging of recombinant phage wherein the vector DNA is cut with one or more restriction enzymes, 3' OH groups of the vector DNA are blocked, vector and DNA to be inserted are contacted under conditions appropriate for ligation of DNA fragments, and the ligation products are
10 tagged with a moiety capable of attaching to reactive 3' OH groups, followed by separation of tagged and untagged molecules.

15 31. A method for at least partially separating nucleic acid molecules in a sample into populations wherein a population is tagged or capable of being tagged with a moiety capable of being immobilised on a matrix either directly or indirectly via a binding partner for the tag, said method comprising contacting the nucleic acid
20 containing sample with a matrix, or, where the tag interacts indirectly with the matrix by means of a binding partner, with a binding partner for the tag and with a matrix, whereby tagged molecules are captured by the matrix and thereby separated from untagged
25 molecules.

32. A method for at least partially separating a mixture of restriction enzyme digested fragments of DNA wherein the starting material is a linear DNA molecule
30 which is tagged or capable of being tagged at or near one or both ends with a moiety capable of being immobilised on a matrix, said method comprising subjecting the DNA molecule to restriction enzyme digestion followed by contacting the sample with a
35 matrix whereby the tagged molecules which originate from an end of the starting material are captured by the matrix and are thereby separated from untagged

09870428 053004

09870123 053001
a 3' primer specific for either the normal or mutant
nucleic acid is used, wherein said primer is
complementary to a region of the nucleic acid where
there is a base difference between the normal and
5 mutated DNA with the 3' terminus of the primer
corresponding to the position in the sample where there
is a difference between the normal and mutant, the
primer being tagged or capable of being tagged, whereby
the presence or absence of the tag in the PCR product is
10 detected.

36. A method of diagnostic PCR in which a test sample
is subjected to PCR with primers complementary to the
normal target DNA wherein the primer for extending at
15 the 3' end of the target anneals to a sequence the 3'
nucleotide of which is mutated in the mutant, the 3'
primer bearing a tag or being capable of being tagged at
or on the 3' nucleotide, wherein the presence of the tag
in the PCR reaction product as detected.

20 37. A method of separating linear from circular nucleic
acid molecules in a sample said method comprising
introducing a tag to an end of a linear nucleic acid
molecule, wherein said tag is capable of being
25 immobilised on a matrix, by direct interaction with the
matrix or by indirect interaction by means of a binding
partner to the tag, and contacting the sample with a
matrix or, where the tag interacts indirectly with the
matrix, with the binding partner to the tag and with a
30 matrix, whereby said tagged linear nucleic acid
molecules are immobilised on the matrix.

38. A method of *in vitro* phage packaging of recombinant
phage wherein the vector DNA is cut with one or more
35 restriction enzymes, 3' OH groups of the vector DNA are
blocked, vector and DNA to be inserted are contacted
under conditions appropriate for ligation of DNA

fragments, and the ligation products are tagged with a moiety capable of attaching to reactive 3' OH groups, followed by separation of tagged and untagged molecules.

09870128 053004